


Ionic Bonding

Giant ionic structures

States of matter



The three states of matter are solid, liquid and gas.

For a substance to change from one state to another, energy must be transferred.

The particles gain energy. This results in the breaking of some of the attractive forces between particles during melting.

To evaporate or boil a liquid, more energy is needed to overcome the remaining chemical bonds between the particles.

Note the difference between boiling and evaporation. When a liquid evaporates, particles leave the surface of the liquid only. When a liquid boils, bubbles of gas form throughout the liquid before rising to the surface and escaping.

The amount of energy needed for a substance to change state is dependent upon the strength of the attractive forces between particles. The stronger the forces of attraction, the more energy needed to break them apart. Substances that have strong attractive forces between particles generally have higher melting and boiling points.

Solid	Liquid	Gas
<p>The particles in a solid are arranged in a regular pattern. The particles in a solid vibrate in a fixed position and are tightly packed together. The particles in a solid have a low amount of kinetic energy.</p> <p>Solids have a fixed shape and are unable to flow like liquids. The particles cannot be compressed because the particles are very close together.</p>	<p>The particles in a liquid are randomly arranged. The particles in a liquid are able to move around each other. The particles in a liquid have a greater amount of kinetic energy than particles in a solid.</p> <p>Liquids are able to flow and can take the shape of the container that they are placed in. As with a solid, liquids cannot be compressed because the particles are close together.</p>	<p>The particles in a gas are randomly arranged. The particles in a gas are able to move around very quickly in all directions. Of the three states of matter, gas particles have the highest amount of kinetic energy.</p> <p>Gases, like liquids, are able to flow and can fill the container that they are placed in. The particles in a gas are far apart from one another which allows the particles to move in any direction.</p> <p>Gases can be compressed; when squashed, the particles have empty space to move into.</p>
Limitations of the Particle Model (HT only)		
<p>The chemical bonds between particles are not represented in the diagrams above.</p> <p>Particles are represented as solid spheres – this is not the case. Particles like atoms are mostly empty space. Particles are not always spherical in nature.</p>		
State Symbols		
<p>In chemical equations, the three states of matter are represented as symbols:</p> <p>solid (s) liquid (l) gas (g) aqueous (aq)</p> <p>Aqueous solutions are those that are formed when a substance is dissolved in water.</p>		
Identifying the Physical State of a Substance		
<p>If the given temperature of a substance is lower than the melting point, the physical state of the substance will be solid.</p> <p>If the given temperature of the substance is between the melting point and boiling point, the substance will be a liquid.</p> <p>If the given temperature of the substance is higher than the boiling point, the substance will be a gas.</p>		

Formation of Ions

Ions are charged particle. They can be either positively or negatively charged, for example Na^+ or Cl^- .

When an element loses or gains electrons, it becomes an ion.

Metals lose electrons to become positively charged.


Non-metals gain electrons to become negatively charged.

Group 1 and 2 elements lose electrons and group 6 and 7 elements gain electrons.

Group	Ions	Element Example
1	+1	$\text{Li} \rightarrow \text{Li}^+ + \text{e}^-$
2	+2	$\text{Ca} \rightarrow \text{Ca}^{2+} + 2\text{e}^-$
6	-2	$\text{Br} + \text{e}^- \rightarrow \text{Br}^-$
7	-1	$\text{O} + 2\text{e}^- \rightarrow \text{O}^{2-}$

Ionic Bonding

Ionic bonding occurs between a metal and a non-metal. Metals lose electrons to become positively charged. Opposite charges are attracted by electrostatic forces – an ionic bond.



Properties of Ionic Compounds

- High melting point – lots of energy needed to overcome the electrostatic forces of attraction.
- High boiling point
- Cannot conduct electricity in a solid as the ions are not free to move.
- Ionic compounds, when molten or in solution, can conduct electricity as the ions are free to move and can carry the electrical current.

Ionic Compounds

Ionic compounds form structures called giant lattices. There are strong electrostatic forces of attraction that act in all directions and act between the oppositely charged ions that make up the giant ionic lattice.

